

Phylogenetic analysis indicates transitions from vegetative to sexual reproduction in the *Lobaria retigera* group (Lecanoromycetidae, Ascomycota)

Carolina CORNEJO, Svetlana CHABANENKO and
Christoph SCHEIDEGGER

Abstract: Phylogenetic relationships among the hypothetical species pairs *Lobaria kurokawae* / *L. retigera* and *L. pseudopulmonaria* / *L. isidiosa* were investigated based on TLC techniques and the phylogenetic analysis of the internal transcribed spacers (ITS) nrDNA. Results of TLC demonstrate that *L. retigera* and *L. kurokawae* can be chemically distinguished from *L. isidiosa* and *L. pseudopulmonaria* by the absence of retigeranic acid. Parsimony analysis of 18 specimens shows two monophyletic clades – a *L. retigera* lineage and a *L. isidiosa* lineage—both including their apotheciate counterparts. Unlike the original hypothesis of species pairs, our study reveals transitions from isidiate morphs that have the potential to rarely form apothecia, towards apotheciate morphs that produce no vegetative propagules.

Key words: evolutionary significant units, lichen-forming ascomycetes, *Lobaria*, species pairs

Introduction

The lichen genus *Lobaria* (Schreb.) Hoffm. includes a number of closely related taxa that are morphologically very similar, except for the production of sexual fruiting bodies or vegetative propagules, and are considered typical examples of species pairs *sensu* Poelt (1970, 1972). Yoshimura (1971) mentioned five species pairs in this genus; two of them in the *Lobaria retigera* group: *L. kurokawae* Yoshim./*L. retigera* (Bory) Trev. and *L. pseudopulmonaria* Gyeln./*L. isidiosa* (Müll. Arg.) Vain. The first species of each pair reproduces sexually with apothecia, while its counterpart propagates mainly asexually with isidia.

The current understanding of the genus *Lobaria* is still controversial. While phylogenetic research based on nuclear and mitochondrial rDNA by Wiklund & Wedin

(2003) demonstrated that *Pseudocyphellaria* is paraphyletic and *Lobaria* formed a monophyletic group, Thomas *et al.* (2002), Stenroos *et al.* (2003) and Miądlikowska & Lutzoni (2004) using nuclear rDNA showed that both *Lobaria* and *Pseudocyphellaria* are polyphyletic. However, these divergent results seem to reflect incongruence between nuclear and mitochondrial DNA that can only be resolved by a multi-gene approach combined with an extensive dataset of *Lobaria* s. lat. Yoshimura (1971) distinguished different sections of *Lobaria* based on ascospore characteristics and the species on further morphological and chemical attributes, but rejected the photobiont as a character of taxonomic value above the species level. Within the *Lobaria retigera* complex he defined species according to chemical characteristics, distinguishing *L. kurokawae* from the morphologically very similar *L. pseudopulmonaria* by the absence of the stictic acid complex. Furthermore, he referred to terpenoids for both taxa but did not identify them. Rao *et al.* (1965, 1966) first described the terpenoids, retigeradiol and retigeranic acid, from samples of the *Lobaria retigera*

C. Cornejo, S. Chabanenko and C. Scheidegger: Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Zuercherstrasse 111, 8903 Birmensdorf, Switzerland. Email: carolina.cornejo@wsl.ch
S. Chabanenko: 2 Sakhalin Botanical Garden, Gorky-street 25, 693023 Yuzhno-Sakhalinsk, Russia.

group but with uncertain taxonomic specifications. Thin layer chromatography (TLC) analysis of Takahashi *et al.* (1972) confirmed the diagnostic system of Yoshimura (1971) distinguishing *L. kurokawae* and *L. retigera* from *L. pseudopulmonaria* and *L. isidiosa* by the absence of stictic acid and the occurrence of retigeric acids A and B. In addition, the authors described a new compound as characteristic for *L. pseudopulmonaria* and *L. isidiosa* that was identified as retigeranic acids A and B by Kaneda *et al.* (1972) and Sugawara *et al.* (1991).

Yoshimura (1971) also considered the production of soredia, isidia or lobules as characteristics for delimiting species; thus, the asexual taxa were treated as different species from their fertile counterparts. Poelt (1970, 1972) and Tehler (1982) understood asexual species as genetically isolated clones because they were considered unable to recombine with sexual relatives. Tehler (1982) also recognized intermediate forms bearing isidia and apothecia that, over time, would lose sexual reproduction completely. However, Mattsson & Lumbsch (1989) argued that sterile species were genetically variable and that it is probably wrong to regard sterile lichens as evolutionary dead ends. The authors further suggested that sterile taxa might have the chance of generating new ones. In fact, all phylogenetic studies on species pairs to date have shown that sterile species are genetically diverse (Lohtander *et al.* 1998*a, b*, 2000; Kroken & Taylor 2001; Myllys *et al.* 1999, 2001; Buschbom & Mueller 2006). In addition, these analyses demonstrated that neither the sexual nor the vegetative reproducing species were monophyletic. Tehler (1982) suggested that the asexual form of species pairs may have developed on multiple occasions from the fertile ones and should only be treated as independent taxa if they have irreversibly lost the ability to produce fruiting bodies. A recent study on evolutionary processes in the species pair *Porpidia flavocoerulescens* and *P. melinodes* may contribute to this debate. Buschbom & Mueller (2006) proposed that the mycobiont might reproduce predominantly with vegetative, symbiotic propagules

until interaction with a more favourable photobiont triggers a switch to the sexual reproductive mode. Consequently, some currently sterile species might gain fertility in a later generation.

In recent years, molecular studies have contributed to our understanding of the phylogenetic relationships between sexual and asexual taxa of lichen forming fungi. The internal transcribed spacers (ITS) of the nuclear rRNA gene cluster have been successfully used in infra-familial and infra-generic reviews (Arup & Grube 2000; Lohtander *et al.* 2000; Ihlen & Ekman 2002; Thomas *et al.* 2002), for delimiting species (Tehler & Källersjö 2001; Miadlikowska *et al.* 2002), in diverse analyses of species pairs (Lohtander *et al.* 1998*a, b*; Kroken & Taylor 2001; Myllys *et al.* 1999, 2001), or in studies of the infraspecific genetic variability (Martín *et al.* 2000; Crespo *et al.* 2002).

The aim of the present study is to examine the phylogenetic relationship between apotheciate and isidiate taxa of the *Lobaria retigera* group. *Lobaria kurokawae* and *L. pseudopulmonaria* consistently form apothecia but never soredia or isidia, whereas *L. retigera* and *L. isidiosa* produce abundant vegetative propagules and rarely form apothecia in addition to the vegetative propagules (Yoshimura 1971). We investigated if the species *L. kurokawae*, *L. retigera*, *L. pseudopulmonaria* and *L. isidiosa* form monophyletic groups. Furthermore, we assessed the taxonomic significance of the chemical compounds by determining whether their differentiation of the two species pairs could be confirmed by the molecular data.

Material and Methods

The specimens studied were collected in Bhutan, Canada, China, Madagascar and Russia. To test the phylogenetic status of the ingroup, we included two species of *Lobaria* from different infrageneric units, *L. amplissima* (Scop.) Forss. and *L. oregana* (Tuck.) Müll. Arg. *Sticta fuliginosa* (Hoffm.) Ach. and *S. canariensis* (Bory) Bory ex Delise were used as outgroup taxa. Information on the taxa, herbarium number and GenBank accession numbers are given in Table 1.

TABLE 1. *Taxa used in this study and the GenBank accession numbers of the ITS nrDNA sequenced for the present study or obtained from GenBank*

Taxon and code		Voucher specimen*	GenBank Accession no.
<i>Lobaria kurokawae</i>	1	Bhutan, Flor Province; GZU 00-411-17/01	DQ419932
	2	China, Yunnan Province; 525SAKH, SCH-CY01	DQ419924
	3	China, Yunnan Province; 548SAKH, SCH-CY14	DQ419942
	4	China, Yunnan Province; 518SAKH, SCH-CY15	DQ419937
	5	Russia, Sakhalin; 498SAKH, SCH-CY18	DQ419938
<i>L. retigera</i>	1	Bhutan, Flor Province; GZU 00-159-10(1)/01	DQ419940
	2	Bhutan, Flor Province; GZU 00-237-10/02	DQ419941
	3	Bhutan, Flor Province; GZU 00-166-34/01	DQ419936
	4	China, Yunnan Province; 544SAKH, SCH-CY52	DQ419930
	5	Canada, British Columbia; SCH-CY09	DQ419934
	6	Canada, British Columbia; SCH-CB05	DQ419929
	7	Russia, Sakhalin; 490SAKH, SCH-CY26	DQ419926
	8	Madagascar; SCH-MP	DQ419928
	9	Madagascar; SCH-X23	EU626996
	10	Madagascar; SCH-X25	EU626997
	11	Madagascar; SCH-X26	EU626998
	12	Madagascar; SCH-X28	EU626999
<i>L. pseudopulmonaria</i>	1	Bhutan, Flor Province; GZU 00-37803/01	DQ419935
	2	Bhutan, Flor Province; GZU 00-183-18/01	DQ419933
	3	Bhutan, Flor Province; GZU 00-339-02/03	DQ419939
	4	China, Yunnan Province; 533SAKH, SCH-CY02	DQ419931
	5	China, Yunnan Province; 530SAKH, SCH-CY53	DQ419925
	6	China, Yunnan Province; 557SAKH, SCH-CY54	DQ419927
<i>L. isidiosa</i>	1	Bhutan, Flor Province; GZU 00-135-18/01	EU627000
	2	China, Yunnan Province; SAKH, SCH-CY30	EU627001
<i>L. oregana</i>		Canada, British Columbia; SCH-CY11	EF605270
<i>Sticta fuliginosa</i>		Canada, British Columbia; SCH-CY32	DQ419943
<i>S. canariensis</i>		Spain, Canary Islands, Tenerife; SCH-X5/01	DQ419944
Sequences obtained from GenBank:			
<i>Lobaria amplissima</i>		Stenroos <i>et al.</i> 2003	AF524925
<i>L. oregana</i>		Goffinet & Goward 1998	AF014111

*Abbreviations for the herbaria. SAKH: Sakhalin Botanical Garden; GZU: Institut für Botanik, Universität Graz; SCH: frozen specimen, collected by C. Scheidegger at WSL.

Chemical analyses

The chemical analyses were performed using standard TLC techniques (Culbertson & Ammann 1979; White & James 1985). Secondary substances were extracted in acetone for approx. 10 min. at 40°C. The extracts were spotted on precoated Merck silica gel F254 plates and eluted in A: toluene : dioxan : acetic acid (180:45:5 ml); B: hexane : diethylether : formic acid (130:100:20 ml); C: toluene : acetic acid (200:30 ml). For each solvent system A, B, and C, we used separate plates with the same extracts. The plates were examined first by UV light at 254 nm and 366 nm wavelengths. The plates were then immersed briefly in a sulphuric acid bath, air dried and heated in an oven at 100°C for approx. 15 min. Identification of the substances was made by comparison with known references from our laboratory. The metabolites of 33 samples were analysed with TLC.

DNA isolation and sequencing

The DNA was isolated using the GenElute Plant Genomic DNA Miniprep Kit (Sigma-Aldrich) according to the manufacture's protocol. The fungal-specific primer ITS1-F (Gardes & Bruns 1993) and the universal primer ITS4 (White *et al.* 1990) were employed to amplify the internal transcribed spacers (ITS 1 and 2) including the 5.8S sequence of the ribosomal RNA gene. The ITS-PCR was performed with 2 µl DNA extract in a total volume of 50 µl containing 1 × PCR buffer (Sigma-Aldrich), 2.0 mM MgCl₂ (Sigma-Aldrich), 40 µM each dNTP (Promega), 80pM each primer, and 0.025 units Taq polymerase (Sigma-Aldrich). The cycling conditions for the ITS region were as follows: a) 2min 94°C, b) 30sec 94°C, c) 30s 57°C, d) 30s 72°C, e) b–d 29 cycles, and f) 10min 72°C.

All PCR products were purified with the QIAquick PCR Purification Kit (QIAGEN), labelled with the Big Dye Terminator v3.1 Kit (Applied Biosystems) and cycle sequenced as follows: 25 cycles of 20s 96°C, 5s 50°C, 2min 60°C. Post-reaction cleanup was carried out in Performa DTR Gel Filtration Cartridges (Edge BioSystems) following the manufacturer's protocol. Sequences were detected in an ABI PRISM 3100 Avant (Applied Biosystems) and checked in the GenBank by nucleotide BLAST searches.

PCR of herbarium specimens

Because several specimens were herbarium samples collected up to ten years ago, the extracted DNA was often of poor quality. PCR amplifications failed in many cases and made the analyses of single-copy genes impossible. Since we had only herbarium samples of the isidiate taxa with retigeranic acid, we applied the following procedures. The ITS PCR was performed as described above and repeated with the first PCR product as template. This method produced the desired PCR fragment but also other artefact fragments. The selected bands were excised from agarose gel (1.5% in TBE Buffer) and purified with the MinElute Gel Extraction Kit (Qiagen) according to the manufacturer's protocol. The cleaned product was used for direct sequencing as described above. The ITS sequences of the specimens *L. isidiota* 1 and 2 were obtained by this technique.

Alignment and phylogenetic analysis

The sequences were assembled pairwise using AutoAssembler 1.4.0 (Applied Biosystems) and optimized by eye. The trimmed sequences were aligned with ClustalX (Thompson *et al.* 1994) with default settings except the following modification of the pairwise and multiple alignment parameters: gap opening 10.00 and gap extension 0.10. In addition, poorly aligned regions were improved using the realign option of ClustalX. The alignment quality was assessed with TuneClustalX (Hall 2004) before and after using modified parameters and the realign option.

Phylogenetic tests were implemented in PAUP* 4.0b10 (Swofford 1998). A heuristic search was carried out with maximum parsimony as criterion and 1000 random-addition-sequence replicates, TBR branch swapping, multrees option in effect, saving all trees and collapsing branches with maximum branch length equal to zero. Gaps were treated as missing data. We set the character polarity by defining in PAUP* *S. fuliginosa* and *S. canariensis* as outgroup. First, we performed two phylogenetic analyses with the unchanged alignment that resulted from ClustalX. Second, ambiguously aligned regions in the ITS sequence were delimited by eye, recoded using the software INAAASE 2.3b according to the author's indications (Lutzoni *et al.* 2000) and re-integrated into the dataset.

The branch support was estimated by bootstrap analysis (Felsenstein 1985) with full heuristic search, 1000 bootstrap replicates, retaining groups with fre-

quency >50%, 100 random-addition-sequences per bootstrap replicate and by saving all trees. Additionally, we calculated Bayesian posterior probabilities with Markov Chain Monte Carlo (MCMC) (Metropolis *et al.* 1953; Hastings 1970) as implemented in MrBayes 3.1 (Huelsenbeck & Ronquist. 2001; Ronquist & Huelsenbeck 2003). Because the substitution model selected by Modeltest (Posada & Crandall 1998) could not be implemented in MrBayes, we executed the analysis with the default substitution model GTR. Nevertheless, the model was simplified assuming equal rates over sites (no gamma variation) to reduce arising burn-out problems (Ronquist *et al.* 2005). The Bayesian analysis was run 115 000 generations, every 10th tree was sampled and the first 2 875 trees were excluded as burn-in.

Reconstructing character evolution

MacClade (Maddison & Maddison 2001) was used to map the occurrence of vegetative propagules and to reconstruct the evolution of this character based on the parsimony criterion and assuming a non-directional evolution. We applied the MP2 tree to infer the reconstruction of the states «with» (0) or «without» (1) vegetative propagules for each taxon. Additionally to these character states, the matrix contained 91 parsimony informative, un-ambiguous nucleotides of the ITS alignment.

Results

Secondary metabolites

All isidiate specimens from Madagascar and one from Bhutan contained abundant retigeric acids A and B (chemotype 1). Other specimens that contained retigeric acid had varying amounts of stictic and norstictic acid (chemotype 2). Finally, chemotype 3 showed copious amounts of the stictic and norstictic acids as well the retigeranic acids A and B. In this chemotype, retigeric acid was not found. If the specimens contained stictic and norstictic acids, the biogenetically related substances, constictic or norconstictic acid, were regularly found. Telephoric acid was consistently found in all specimens and was therefore not used for characterization of the chemotypes. The results of TLC analyses are summarized in Fig. 1.

ITS analysis

A total of 28 sequences was produced, but not all were used for phylogenetic analysis. Nevertheless, all sequences were submitted





FIG. 1. Summary of chemical patterns found in TLC analysis of *Lobaria retigera* group. Chemotypes 1 and 2 are characteristic of the *L. retigera* lineage and chemotype 3 of the *L. isidiosa* lineage. Lichens which contain large amounts of retigeric acid (large circles) were observed to produce less (small circles, chemotype 2) or no stictic acid.

to GenBank (Table 1). The dataset included 23 sequences of apotheciate taxa and their counterpart species as well as *L. oregana*, *L. amplissima* and the two outgroup taxa. After sequences were trimmed so that they had a concerted beginning and end point, the dataset consisted of 476 nucleotide sites. The first heuristic search was performed in PAUP* with maximum parsimony including all nucleotides (MP1). PAUP* defined gaps as missing, excluded 330 constant sites and carried out the heuristic search with 112 nucleotides that were parsimoniously informative. This analysis produced one most parsimonious tree of 255 steps length with the tree score indices CI=0.773 and RI=0.841. Second, for the MP2 analysis 385 gapped, ambiguously aligned, constant or parsimony uninformative sites were excluded as performed by the command include/exclude characters in PAUP* and the remaining 91 informative sites resulted in two

phylogenetic trees of 164 steps, CI=0.726 and RI=0.845. Finally, for the MP3 analysis, a total of 91 nucleotides, located in 13 ambiguously aligned regions, were excluded and replaced by 13 recoded characters (INAASE). Further, 324 gapped, constant or uninformative nucleotides were excluded, thus 74 parsimony informative characters could be integrated into the heuristic search. The MP3 analysis revealed 47 most parsimonious trees of 167 steps, CI=0.802 and RI=0.866. All three analyses produced phylogenetic trees with very similar structure except for some terminal branches. The bootstrap cladograms implemented in PAUP* did not resolve all branches of the phylogenetic trees and not all nodes obtained enough bootstrap support (>75%). The resulting phylogenetic tree obtained with MrBayes was very similar to the MP phylograms. However, the posterior probabilities tree positioned the *L. oregana*-*L. amplissima* branch out of the ingroup. Consequently, no posterior probability was obtained for this node. As there were no contradictions for the ingroup studied among the MP1, MP2, MP3 trees and the Bayesian phylogram, the MP2 tree with fewest substitutions (164 steps) was chosen for presentation (Fig. 2).

Outgroup rooting in the MP2 analysis led to a cluster with *L. oregana* and *L. amplissima* basal to the ingroup in question. The ingroup was maximally supported ($b=100$; $MrB=100$) and revealed two lineages. Each lineage included both apotheciate and isidiate specimens with high bootstrap support ($b=89$; $MrB=100$ and $b=82$; $MrB=100$), but within each lineage, apotheciate and isidiate taxa were intermingled. The order of taxa in the first clade indicated in some cases geographical patterns, for example, Malagasy ($b=96$; $MrB=100$) or Canadian specimens ($b=86$; $MrB=100$) fit together with high bootstrap support. This was not found in the *L. isidiosa* lineage. The MP2 position of the Sakhalin specimens was supported neither by bootstrap analysis nor by posterior probabilities. In the posterior probability tree these specimens formed one clade together with the Malagasy samples, although with minor support ($MrB=71$).



Discussion

We did not examine any type material in this study and as only a limited number of recently collected specimens was available for molecular studies, our data do not allow us to draw nomenclatural conclusions from the chemical and molecular investigations. The oldest name at the species level in the *L. retigera* group is *L. retigera*, described from La Réunion (Bory 1804; see Yoshimura 1971 for a nomenclatural discussion). It is likely that the name *L. retigera* applies to the specimens belonging to the retigeric acid chemotype from Madagascar. The name *L. kurokawae* is widely used in literature for apotheciate specimens but would be a younger facultative synonym of *L. retigera* if the African, Asian and American populations of the retigeric acid chemotype were to be considered conspecific. In this study, *L. kurokawae* is used for apotheciate specimens that produce retigeric acid but chemical analyses should reveal if facultative synonyms of *L. retigera* would have nomenclatural priority if additional molecular analyses revealed a regional subdivision of the chemotypes 1 and 2. For chemotype 3, *L. isidiosia* would have nomenclatural priority and in our study the name *L. pseudopulmonaria* is solely used to indicate apotheciate specimens that belong to chemotype 3.

The presence or absence of stictic or norstictic acids showed no relation to the clades found in the ITS phylogeny. While the chemical pattern of the *L. retigera* lineage is characterized by the production of retigeric acid (chemotypes 1 and 2), TLC analyses indicate that *L. pseudopulmonaria* and *L. isidiosia* differ from the other *Lobaria* species studied by the presence of retigeranic acid (chemotype 3). In addition, we observed that lichens of the chemotypes 1 and 2, which contain

large amounts of retigeric acid, have less or no stictic acid indicating a negative correlation between these metabolites (Fig. 1).

On the basis of the presence of retigeric or retigeranic acids and on the ITS phylogeny, two lineages showing a high bootstrap support were distinguished (Fig. 2). One lineage includes the apotheciate taxon *L. kurokawae* and the isidiate taxon *L. retigera* and the second lineage contains the apotheciate *L. pseudopulmonaria* and the isidiate *L. isidiosia*. In addition, the phylogenetic tree shows that the apotheciate and the isidiate forms within each lineage are not monophyletic. Highly supported groups within the *L. retigera* lineage are formed among specimens originating from the same geographical region but from different localities. This was found among specimens where only isidiate forms were available from regions such as Madagascar and the forests of British Columbia, but the same pattern was also found if the specimens showed different reproduction modes. For example, the apotheciate and isidiate specimens from Yunnan form a highly supported clade. These results suggest a strong geographical structure in the *L. retigera* lineage, among the populations in Canada, Madagascar, Bhutan, Yunnan (China), and Sakhalin (Russia), of this widely distributed taxon. It is likely that these isolated populations have evolved over geologically long periods, and, independent of their reproductive mode and taxonomic rank, these geographically differentiated clades may deserve the status of evolutionarily significant units (Moritz 1995) for biological conservation. No comparable levels of genetic differentiation have so far been found in the *L. isidiosia* clade where the samples originated from a more restricted area (Bhutan and Yunnan).

The evolution of the character, vegetative propagules, shows an equivocal presence/

FIG. 2. One of two most parsimonious trees produced by phylogenetic analysis of the ITS 1 and 2 sequences and the 5.8S gene showing the relationship of the isidiate species *L. retigera* and *L. isidiosia* and their apotheciate counterparts *L. kurokawae* and *L. pseudopulmonaria*. *Sticta canariensis* and *S. fuliginosa* were defined as the outgroup. ○, chemotype 1+2: retigeric acids A & B + stictic and norstictic acids; ●, chemotype 3: retigeranic acids A & B + stictic and norstictic acids. Numbers beside internodes represent the percentage of 1000 bootstrap replicates (left) and the Bayesian posterior probabilities (right). Only values $\geq 50\%$ are shown; CI: 0.726, RI: 0.845. Species names in bold type with vegetative propagules; species names in normal type without vegetative propagules.

TABLE 2. Diagnostic characteristics of *Lobaria* species involved in this study

Taxon	<i>L. kurokawae</i> / <i>L. retigera</i> apotheciate/isidiate		<i>L. pseudopulmonaria</i> / <i>L. isidiosa</i> apotheciate/isidiate	
	(1)	Chemotype 1+2 (2)	(1)	Chemotype 3 (2)
Stictic and/or norstictic acid	–	–/+ / ++	++	++
Retigeric acids A and B	++	+ / ++	–	–
Retigeranic acids A and B	–	–	+	++

(1) Takahashi *et al.* (1972); (2) results of this study – Not found; + traces; ++ abundant

absence of isidia at the base of the *L. isidiosa* lineage. A more intensive taxon sampling will be needed to identify the ancestral character state in this clade. However, in the *L. retigera* lineage the internal nodes were revealed to be isidiate and only three terminal nodes developed into the non-isidiate character state that is typical for the apotheciate taxon. On the other hand, morphs of *L. isidiosa* and *L. retigera* bearing both isidia and apothecia (Yoshimura 1971) show that sexual reproduction of thalli is suppressed rather than lost and hence recombination between apotheciate and isidiate lineages can be expected. Recently, Buschbom & Mueller (2006) showed in *Porpidia flavocoerulescens* and *P. melinodes* that the transition from the vegetative to the sexual state happens more frequently than vice versa which implies that the isidiate morphs may not have completely lost the ability to build fruiting bodies. All studies to date, including the present study, have demonstrated that sexually and vegetative reproducing species pairs are not monophyletic and, consequently, question the hypothesis of genetic isolation of vegetative lineages. Moreover, unlike other studies where the sorediate specimens supposedly arose several times at different geographical locations from fertile lineages (Lohtander *et al.* 1998a, b; Myllys *et al.* 1999; Myllys *et al.* 2001), our study of *L. retigera* shows that apotheciate clades have developed repeatedly from rarely fertile, isidiate morphs.

In contrast to Poelt's (1972) hypothesis of species pairs, published studies reveal transitions from isidiate morphs that have the potential to form apothecia rarely, towards

apotheciate morphs that produce no vegetative propagules. The widespread green-algal *Lobaria pulmonaria* has a combined reproductive strategy, i.e., it always produces soredia but only rarely forms apothecia. Zoller *et al.* (1999) suggested that *L. pulmonaria* is heterothallic and is likely to remain sterile in populations with low genetic diversity. Species with infrequent formation of apothecia were often shown to be heterothallic (Zoller *et al.* 1999; Kroken & Taylor 2001; Scherrer *et al.* 2004; Seymour *et al.* 2005; Honegger & Zippler 2007). Because several examples of abundantly fertile heterothallic species such as *Ramalina fastigiata*, *R. fraxinea*, *Physcia aipolia* and *P. stellaris* (Honegger & Zippler 2007) are known, it remains unclear if a change in the frequency of apothecia is paralleled by a transition from heterothallism to homothallism within a lichen species such as *L. retigera*.

We are very grateful to W. Obermayer (Graz) and C. Keller (Birmensdorf) for collaborating on chemical analysis and U. Kamm (Birmensdorf) for providing the specimen from Tenerife. We wish to thank S. Werth (Birmensdorf) and the anonymous referees for constructive comments. S. Chabanenko and C. Scheidegger were supported by a research grant from the Swiss National Science Foundation (Scopes 7SUPJ062353 and IB73AO-111137/1).

REFERENCES

- Arup, U. & Grube, M. (2000) Is *Rhizoplaca* (Lecanorales, lichenized Ascomycota) a monophyletic genus? *Canadian Journal of Botany* **78**: 318–327.
- Bory de Saint-Vincent, J. B. G. M. (1804) *Voyage dans les Quatre Principales Îles des Mers d'Afrique*. Vols 1–3. Paris.
- Buschbom, H. & Mueller, G. M. (2006) Testing “species pair” hypotheses: evolutionary processes in

- the lichen-forming species complex *Porpidia flavo-coerulescens* and *Porpidia melinodes*. *Molecular Biology and Evolution* **23**: 574–586.
- Crespo, A., Molina, M. C., Blanco, O., Schroeter, B., Sancho, L. G. & Hawksworth, D. L. (2002) rDNA ITS and β -tubulin gene sequence analyses reveal two monophyletic groups within the cosmopolitan lichen *Parmelia saxatilis*. *Mycological Research* **106**: 788–795.
- Culberson, C. F. & Ammann, K. (1979) Standard-methode zur Dünnschichtchromatographie von Flechtensubstanzen. *Herzogia* **5**: 1–24.
- Felsenstein, J. (1985) Confidence-limits on phylogenies: an approach using the bootstrap. *Evolution* **39**: 783–791.
- Gardes, M. & Bruns, T. D. (1993) ITS primers with enhanced specificity for Basidiomycetes—application to the identification of mycorrhizae and rusts. *Molecular Ecology* **2**: 113–118.
- Goffinet, B. & Goward, T. (1998) Is *Nephroma silvae-veteris* the cyanomorph of *Lobaria oregana*? Insights from molecular, chemical and morphological characters. In *Lichenographia Thomsoniana: North America Lichenology in Honour of John W. Thomson* (M. G. Glenn, R. C. Harris, R. Dirig & M. S. Cole, eds): 41–52. Ithaca, NY: Mycotaxon Ltd.
- Hall, B. G. (2004) *Improving ClustalX Alignments with TuneClustalX*. Bellingham Research Institute Report No. 1. www.bellinghamresearchinstitute.com
- Hastings, W. K. (1970) Monte Carlo sampling methods using Markov chains and their applications. *Biometrika* **57**: 97–109.
- Honegger, R. & Zippler, U. (2007) Mating systems in representatives of Parmeliaceae, Ramalinaceae and Physciaceae (Lecanoromycetes, lichen-forming ascomycetes). *Mycological Research* **111**: 424–432.
- Huelsenbeck, J. P. & Ronquist, F. (2001) MrBayes: Bayesian inference of phylogeny. *Bioinformatics* **17**: 754–755.
- Ihlen, P. G. & Ekman, S. (2002) Outline of phylogeny and character evolution in *Rhizocarpon* (Rhizocarpaceae, lichenized Ascomycota) based on nuclear ITS and mitochondrial SSU ribosomal DNA sequences. *Biological Journal of the Linnean Society* **77**: 535–546.
- Kaneda, M., Takahashi, R., Iitaka, Y. & Shibata, S. (1972) Retigeranic acid, a novel sesterpene isolated from the lichens of *Lobaria retigera* group. *Tetrahedron Letters* **45**: 4609–4611.
- Kroken, S. & Taylor, J. W. (2001) Outcrossing and recombination in the lichenized fungus *Letharia*. *Fungal Genetics and Biology* **34**: 83–92.
- Lohtander, K., Källersjö, M. & Tehler, A. (1998a) Dispersal strategies in *Roccellina capensis* (Arthoniales). *Lichenologist* **30**: 341–350.
- Lohtander, K., Myllys, L., Sundin, R., Källersjö, M. & Tehler, A. (1998b) The species pair concept in the lichen *Dendrographa leucophaea* (Arthoniales): analyses based on ITS sequences. *Bryologist* **101**: 404–411.
- Lohtander, K., Källersjö, M., Moberg, R. & Tehler, A. (2000) The family Physciaceae in Fennoscandia: phylogeny inferred from ITS sequences. *Mycologia* **92**: 728–735.
- Lutzoni, F., Wagner, P., Reeb, V. & Zoller, S. (2000) Integrating ambiguously aligned regions of DNA sequences in phylogenetic analyses without violating positional homology. *Systematic Biology* **49**: 628–651.
- Maddison, D. E. & Maddison, W. P. (2001) *MacClade 4: Analysis of Phylogeny and Character Evolution. Version 4.03*. Sunderland, Mass.: Sinauer Associates.
- Martín, M. P., Winka, K., Llimona, X. & Lumbsch, H. T. (2000) Evaluation of morphological variation in the lichen *Diploschistes ocellatus* (Ascomycota, Ostropales): evidence from nuclear rDNA ITS sequence data. *Plant Biology* **2**: 571–578.
- Mattsson, J. E. & Lumbsch, H. T. (1989) The use of the species pair concept in lichen taxonomy. *Taxon* **38**: 238–241.
- Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H. & Teller, E. (1953) Equations of state calculations by fast computing machines. *Journal of Chemical Physics* **21**: 1087–1091.
- Miadlikowska, J., McCune, B. & Lutzoni, F. (2002) *Pseudocyphellaria perpetua*, a new lichen from Western North America. *Bryologist* **105**: 1–10.
- Miadlikowska, J. & Lutzoni, F. (2004) Phylogenetic classification of Peltigeralean fungi (Peltigerales, Ascomycota) based on ribosomal RNA small and large subunits. *American Journal of Botany* **91**: 449–464.
- Moritz, C. (1995) Uses of molecular phylogenies for conservation. *Philosophical Transactions of the Royal Society of London, Series B* **349**: 113–118.
- Myllys, L., Lohtander, K., Källersjö, M. & Tehler, A. (1999) Sequence insertions and ITS data provide congruent information on *Rocella canariensis* and *R. tuberculata* (Arthoniales, Euascomycetes) phylogeny. *Molecular Phylogenetics and Evolution* **12**: 295–309.
- Myllys, L., Lohtander, K. & Tehler, A. (2001) β -tubulin, ITS and group I intron sequences challenge the species pair concept in *Physcia aipolia* and *P. caesia*. *Mycologia* **93**: 335–343.
- Poelt, J. (1970) Das Konzept der Artenpaare bei den Flechten. In *Flechtensymposium 1969, Deutsche Botanische Gesellschaft, 1970, Stuttgart, Germany*.
- Poelt, J. (1972) Die taxonomische Behandlung von Artenpaaren bei den Flechten. *Botaniska Notiser* **125**: 77–81.
- Posada, D. & Crandall, K. A. (1998) Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**: 817–818.
- Rao, P. S., Sarma, K. G. & Seshadri, T. R. (1965) Chemical components of *Lobaria* lichens from Western Himalayas. *Current Science* **34**: 9–11.
- Rao, P. S., Sarma, K. G. & Seshadri, T. R. (1966) Chemical components of *Lobaria subsidiosa*, *L. retigera* and *L. subretigera* from Western Himalayas. *Current Science* **35**: 147–148.

- Ronquist, F. & Huelsenbeck, J. P. (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572–1574.
- Ronquist, F., Huelsenbeck, J. P. & van der Mark, P. (2005) *Mr Bayes 3.1 Manual. Draft 5/26/2005*. <http://mr bayes.csit.fsu.edu/manual.php>
- Scherrer, S., Zippler, U. & Honegger, R. (2004) Characterisation of the mating-type locus in the genus *Xanthoria* (lichen-forming ascomycetes, lecanoromycetes). *Fungal Genetics and Biology* **42**: 976–988.
- Seymour, F. A., Crittenden, P. D., Dickinson, M. J., Paoletti, M., Montiel, D., Cho, L. & Dyer, P. S. (2005) Breeding systems in the lichen-forming fungal genus *Cladonia*. *Fungal Genetics and Biology* **42**: 554–563.
- Stenroos, S., Stocker-Wörgötter, E., Yoshimura, I., Myllys, L., Thell, A. & Hyvonen, J. (2003) Culture experiments and DNA sequence data confirm the identity of *Lobaria* photomorphs. *Canadian Journal of Botany* **81**: 232–247.
- Sugawara, H., Kasuya, A., Iitaka, Y. & Shibata, S. (1991) Further-studies on the structure of retigeranic acid. *Chemical and Pharmaceutical Bulletin* **39**: 3051–3054.
- Swofford, D. L. (1998) *PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods)*. Version 4. Sinauer Associates, Sunderland, Massachusetts.
- Takahashi, R., Chiang, H. C., Aimi, N., Tanaka, A. & Shibata, S. (1972) The structures of retigeric acids A and B from lichens of the *Lobaria retigera* group. *Phytochemistry* **11**: 2039–2045.
- Tehler, A. (1982) The species pair concept in lichenology. *Taxon* **31**: 708–714.
- Tehler, A. & Källersjö, M. (2001) *Parmeliopsis ambigua* and *P. hyperopta* (Parmeliaceae): species or chemotypes? *Lichenologist* **35**: 403–408.
- Thomas, M. A., Ryan, D. J., Farnden, K. J. F. & Galloway, D. J. (2002) Observations on phylogenetic relationships within Lobariaceae Chevall. (Lecanorales, Ascomycota) in New Zealand, based on ITS-5.8S molecular sequence data. *Bibliotheca Lichenologica* **82**: 123–138.
- Thompson, J. D., Higgins, D. G. & Gibson, T. J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Research* **22**: 4673–4680.
- White, F. J. & James, P. W. (1985). *A New Guide to Microchemical Techniques for the Identification of Lichen Substances*. London: British Lichen Society.
- White, T. J., Bruns, T., Lee, S. & Taylor, J. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In *PCR Protocols: a Guide to Methods and Applications* (M. A. Innis, D. H. Gelfond, J. J. Sninsky & T. J. White, eds): 315–322. New York: Academic Press, Inc.
- Wiklund, E. & Wedin, M. (2003) The phylogenetic relationships of the cyanobacterial lichens in the Lecanorales suborder Peltigerineae. *Cladistics* **19**: 419–431.
- Yoshimura, I. (1971) The genus *Lobaria* of Eastern Asia. *Journal of the Hattori Botanical Laboratory* **34**: 231–336.
- Zoller, S., Lutzoni, F. & Scheidegger, C. (1999) Genetic variation within and among populations of the threatened lichen *Lobaria pulmonaria* in Switzerland and implications for its conservation. *Molecular Ecology* **8**: 2049–2059.

Accepted for publication 12 January 2009